

‘ABSTRACT’ HOMOMORPHISMS OF l -ADIC GROUPS AND ABELIAN VARIETIES

SIMAN WONG FEBRUARY 1, 2008 – 07:44 **DRAFT**

ABSTRACT. Let k be a totally real field, and let A/k be an absolutely irreducible, polarized Abelian variety of odd, prime dimension whose endomorphism rings is non-trivial and is defined over k . Then the only strictly compatible families of abstract, absolutely irreducible representations of $\mathrm{Gal}(\bar{k}/k)$ coming from A are tensor products of Tate twists of symmetric powers of two-dimensional λ -adic representations plus field automorphisms. The main ingredients of the proofs are the work of Borel and Tits on the ‘abstract’ homomorphisms of almost simple algebraic groups, plus the work of Shimura on the fields of moduli of Abelian varieties.

1. INTRODUCTION

Let A be an absolutely irreducible Abelian threefold over a number field k . Suppose that the endomorphisms of A are all defined over k . Let l be a prime, and write $T_l(A)$ for the l -adic Tate module of A . Let $L/\mathbf{Q}_l$ be a finite extension, set $G_k := \mathrm{Gal}(\bar{k}/k)$, and let $\rho : G_k \rightarrow GL_3(L)$ be an *abstract* (i.e. not necessarily continuous) group homomorphism which is a quotient of the l -adic representation $G_k \rightarrow \mathrm{Aut}(T_l(A))$ by an abstract homomorphism $\mathrm{Aut}(T_l(A)) \rightarrow GL_3(L)$. A special case of our main results (Theorems 2 and 3) says that if l is sufficiently large and if k is *totally real*, then ρ is the Tate twist of the symmetric-square of an l -adic representation $G_k \rightarrow GL_2(L)$ plus a field automorphism $L \rightarrow L$. To put this result in perspective, we first recall how l -adic representations arise in the Langlands program.

Denote by $\mathbb{A}_k$ the adeles of a number field k . The Langlands program predicts that algebraic cuspidal automorphic forms on $GL_n(\mathbb{A}_k)$ correspond to motives, or, more concretely, compatible families of n -dimension l -adic representations coming from étale cohomology of algebraic varieties defined over k . For *holomorphic* eigenforms on $GL_2(\mathbb{A}_{\mathbf{Q}})$ this conjecture follows from the work of Eichler-Shimura, Deligne, and Deligne-Serre, by relating the eigenforms to the cohomology of certain Kuga fiber varieties. This has been extended to holomorphic GL_2 -eigenforms over totally real fields, by Blasius-Rogawski [5] and Taylor [27]. When the automorphic form corresponds to a selfdual cuspidal representation, Clozel establishes the existence of these l -adic representations under mild conditions [8, p. 150]. Beyond that little is known for non-selfdual representations for $n \geq 3$.

Clozel raises the question of whether we can collect numerical evidence to support the Langlands correspondence for non-selfdual representations. Since then several authors have carried out numerical studies in the smallest open case $n = 3$ ([1], [3], [11], [12]). Such investigations have two parts. On the automorphic side, there is a well-known connection between cuspidal automorphic forms on $GL_3(\mathbb{A}_{\mathbf{Q}})$ and cuspidal cohomology of arithmetic

1991 *Mathematics Subject Classification.* Primary 11G5, 11G10; Secondary 11G15.

Key words and phrases. Abelian varieties, endomorphisms, Galois representations, moduli.

subgroup of $SL_3(\mathbf{Z})$ (cf. [17] for an exposition). Thus automorphic forms on $GL(3)$ can be systematically enumerated, at least for small levels, via Ash's modular symbol algorithm on $GL(3)$ [2]. On the Galois side, however, there is no known algorithm that generates all irreducible 3-dimensional l -adic representations. Examples in the literature come from either ingenious but ad hoc geometric constructions ([11], [12]), or, in the case of mod- l representations, number fields with special Galois groups ([1], [3]).

A familiar source of compatible families of l -adic representations comes from the Tate modules of Abelian varieties. It is natural to try to extract from them non-selfdual representations. To be specific, denote by F_λ the completion of a number field F at a finite place λ of residual characteristic l . Suppose an abstract homomorphism¹ $\rho_\lambda : G_k \rightarrow GL_N(F_\lambda)$ comes from A/k , i.e. there exists an abstract homomorphism $\iota_\lambda : \text{Aut}(T_l(A)) \rightarrow GL_N(\overline{F_\lambda})$ such that, along with the canonical inclusion

$$j_\lambda : GL_N(F_\lambda) \hookrightarrow GL_N(\overline{F_\lambda}),$$

the following diagram commutes ($V_l(A) := T_l(A) \otimes \mathbf{Q}_l$):

$$(1) \quad \begin{array}{ccccc} G_k & \xrightarrow{\rho_{A,l}} & \text{Aut}(T_l(A)) & \subset & \text{Aut}(V_l(A)) \\ \rho_\lambda \downarrow & & & & \downarrow \iota_\lambda \\ GL_N(F_\lambda) & \xrightarrow{j_\lambda} & GL_N(\overline{F_\lambda}) & & \end{array}$$

Suppose A/k is an Abelian threefold such that $\text{End}(A/k) \otimes \mathbf{Q}$ contains a quadratic field. Then $V_l(A)$ splits into a direct sum of two conjugate, odd-dimensional Galois representations. Furthermore, if $\text{End}(A/\overline{k}) \otimes \mathbf{Q}$ is exactly a quadratic field, by the Mumford-Tate conjecture [19] we expect the Lie algebra of the image of these conjugate representations to contain $\mathfrak{sl}_3(\mathbf{Z}_l)$. The Jacobian of a generic Picard quartic $y^3 = x^4 + ax^3 + bx^2 + cx + d$ over k would satisfy the required conditions, thereby yielding interesting irreducible, non-selfdual 3-dimensional l -adic representations, *provided that* k contains a primitive third-root of unity. One important consequence of the result stated in the first paragraph is that such examples cannot exist if k is not totally real, even if we do not require that the representation be continuous.

The proofs of our main results have two parts. As a preliminary step, we utilize Shimura's work on the fields of moduli of Abelian varieties with PEL structures to limit the possibilities for the endomorphism algebras of odd-dimensional Abelian varieties over a *totally real* field.

Theorem 1. *Let A be an absolutely simple, polarized Abelian variety of odd dimension d defined over a totally real number field k . Suppose that the endomorphisms of A are all defined over k . Then $E := \text{End}(A/k) \otimes \mathbf{Q}$ is a totally real number field of degree dividing d .*

With A/k as in Theorem 1, set $[E : \mathbf{Q}] = g$ and $\delta = d/g$. Since the action of E on $V_l(A)$ commutes that of G_k , we have a direct sum decomposition

$$(2) \quad \rho_{A,l} \simeq \pi_{l,1} \oplus \cdots \oplus \pi_{l,g},$$

¹we will explain shortly will we want to work with not-necessarily continuous maps.

where each $\tau_{l,m}$ is a representation $G_k \rightarrow GL_{2\delta}(E_{l,m})$ for some completion $E_{l,m}$ of E at a place above l . Denote by

$$\pi_{l,m} : \text{Aut}(T_l(A)) \rightarrow GL_{2\delta}(E_{l,m})$$

the corresponding quotient. If ρ_λ is irreducible, it must factor through one of the $\tau_{l,m}$. If ι_λ is further assumed to be l -adic analytic, standard Lie theory then implies that ι_λ is the composition of some $\pi_{l,m}$ with a rational² representation $GL_{2\delta}(E_{l,m}) \rightarrow GL_N(\overline{F}_\lambda)$. However, if we start out with λ -adic representations that arise in arithmetic geometry, we only know that ρ_λ is continuous. So the additional hypothesis that ρ_λ comes from an Abelian variety does not automatically imply that ι_λ is continuous, let alone being λ -adic analytic. To address this issue, we invoke a deep result of Borel and Tits [6] which asserts that, over an infinite field K , the ‘abstract’ homomorphisms of an almost simple matrix group G over K into $PGL_n(\overline{K})$ are tensor products of the rational projective representations of G plus field automorphisms; additional work is required to handle ordinary representations. Before we state our main results we introduce some notation.

Let V be a two-dimension vector space over a field K . Let $\varphi : K \rightarrow \overline{K}$ be a field homomorphism. For any integer $n \geq 1$, denote by $\sigma_\varphi^{(n)} \circ \varphi$ the n -th symmetric power of the ‘ φ -twisted’ standard representation $GL_2(K) \rightarrow GL_2(\varphi(K)) \subset GL_2(\overline{K})$. This is a $(n+1)$ -dimensional representation.

Theorem 2. *With the notation as in Theorem 1, suppose further that there exists an abstract, absolutely irreducible representation $\rho_\lambda : G_k \rightarrow GL_N(F_\lambda)$ coming from A . Suppose $l = \text{char}(\lambda)$ is sufficiently large.*

(a) *If $N < 2d$ and ρ_λ is non-Abelian then $[E : \mathbf{Q}] > 1$.*

(b) *Suppose $d = [E : \mathbf{Q}]$. Then there exist*

- *a choice $\tau_{l,m}$ among the factors in the decomposition (2),*
- *positive integers $r(\lambda)$ and $n_1(\lambda), \dots, n_{r(\lambda)}(\lambda)$ with $\sum_{i=1}^{r(\lambda)} (n_i(\lambda) + 1) = N$,*
- *a field homomorphism $\varphi_i(\lambda) : E_{l,m} \rightarrow \overline{E}_{l,m}$ for each $1 \leq i \leq r(\lambda)$, and*
- *a homomorphism of multiplicative groups $f_\lambda : E_{l,m}^\times \rightarrow \overline{E}_{l,m}^\times$,*

such that, in the notation of diagram (1),

$$\iota_\lambda = \left(\left[\bigotimes_{i=1}^{r(\lambda)} \sigma_{\varphi_i(\lambda)}^{(n_i(\lambda))} \circ \varphi_i(\lambda) \right] \otimes (f_\lambda \circ \det) \right) \circ \tau_{l,m}.$$

Remark 1. (a) Set $d = N = 3$ and we recover the result stated in the first paragraph.

(b) Our technique applies *mutatis mutandis* when $[E : \mathbf{Q}] < d$, once we replace the symmetric power representations of GL_2 by the Weyl modules associated to GL_n [10, §15.5]. The same goes for Theorem 3 below. In addition, Theorems 2 and 3 remain true if we replace λ -adic representations by mod- λ representations.

In practice, continuous λ -adic representations often arise in a compatible family. Our next result says that if $\{\rho_\lambda\}_\lambda$ is a compatible family coming from an Abelian variety A/k as in Theorem 1, then the parameters $r(\lambda), n_i(\lambda), f_\lambda$ and $\varphi_i(\lambda)$ in Theorem 2 are essentially

²i.e. a morphism of algebraic groups.

independent of λ . To set up the notation we first review the definition of compatible families of representations.

Definition. A strictly compatible family of λ -adic rational representations of G_k consists of

- a number field $F/\mathbf{Q}$,
- a collection $\{\rho_\lambda : G_k \rightarrow GL_n(F_\lambda)\}_\lambda$ of continuous Galois representations for each finite place λ of F ,
- a finite set S of finite places of $\mathcal{O}_k$,

such that, with $\Phi_\lambda : F \rightarrow F_\lambda$ denoting the canonical embedding,

- for any prime $\mathfrak{p} \notin S \cup \{\lambda\}$, the coefficients of the characteristic polynomial $\text{char}(\rho_\lambda, \mathfrak{p})$ of $\rho_\lambda(\text{Frob}_\mathfrak{p})$ lies in $\Phi_\lambda(F)$, and
- for any two distinct finite places $\lambda, \mathfrak{q}$ and for any place $\mathfrak{p} \notin S \cup \{\lambda, \mathfrak{q}\}$,

$$\Phi_\lambda^{-1}(\text{char}(\rho_\lambda, \mathfrak{p})) = \Phi_\mathfrak{q}^{-1}(\text{char}(\rho_\mathfrak{q}, \mathfrak{p})).$$

Remark 2. The phrase ‘rational representations’ refers to the condition that the image of each ρ_λ lies in $\Phi_\lambda(F)$ for a fixed number field F , rather than the whole $\overline{F}_\lambda$. In particular, this is different from the terminology in the theory of algebraic groups (cf. footnote 2).

Theorem 3. Let A/k be as in Theorem 1 with $[E : \mathbf{Q}] = d$. Suppose $\{\rho_\lambda\}_\lambda$ is a strictly compatible family of absolutely irreducible rational representations coming from A/k . Following the notation in Theorem 2, for all but finitely many pairs $(\lambda, \mathfrak{q})$ of finite places of k ,

- (i) $r(\lambda) = r(\mathfrak{q})$,
- (ii) $n_i(\lambda) = n_i(\mathfrak{q})$ for all i , up to reordering,
- (iii) $f_\lambda|_{\mathbf{Z}} = f_\mathfrak{q}|_{\mathbf{Z}}$, and
- (iv) there exists a field isomorphism $\xi_{\lambda, \mathfrak{q}} : \overline{\Phi_\lambda(E)} \rightarrow \overline{\Phi_\mathfrak{q}(E)}$ such that

$$\xi_{\lambda, \mathfrak{q}}(\varphi_i(\lambda)|_{\Phi_\lambda(E)}) = (\varphi_i(\mathfrak{q})|_{\Phi_\mathfrak{q}(E)}) \quad \text{for every } 1 \leq i \leq r(\lambda).$$

Remark 3. We clarify the meaning of conditions (iii) and (iv). In the notation of Theorem 2 (and dropping the λ from $r(\lambda)$ and $n_i(\lambda)$),

$$\iota_\lambda = \left(\left[\bigotimes_{i=1}^r \sigma_{\varphi_i}^{(n_i)} \circ \varphi_i(\lambda) \right] \otimes (f_\lambda \circ \det) \right) \circ \tau_{l, m}.$$

Since $\Phi_\lambda(E) \subset E_\lambda$ is dense in E_λ , for the purpose of computing ι_λ , condition (iv) completely determines φ_λ . Similarly, for A/k as in Theorem 1, $\det \tau_{l, m}$ takes values in $\mathbf{Z}_l - \{0\}$ and $\mathbf{Z}$ is dense in $\mathbf{Z}_l$, so condition (iii) completely determines f_λ .

Since G_k is compact, if $\rho_\lambda : G_k \rightarrow GL_N(F_\lambda)$ is continuous then its image is conjugate to a subgroup of $GL_N(\mathcal{O}_\lambda)$, where $\mathcal{O}_\lambda$ denotes the ring of integers of F_λ (cf. [18, Remark 1 on p. 1]). To say that ρ_λ comes from A/k is to say that the $\mathcal{O}_\lambda[G_k]$ -module for ρ_λ , *after tensoring with $\mathbf{Q}$* , is a quotient of $V_l(A)$. This suggests that we seek an *integral* form of our results, i.e. *without* taking the tensor product. This calls for an integral form of the work of Borel-Tits, i.e. a characterization of abstract homomorphisms of linear groups such as

$GL_n(\mathcal{O}_\lambda)$ into $GL_m(\mathcal{O}'_\lambda)$. This is a very interesting question by itself, but little is known in this direction. For the record we have the following partial results.

Theorem 4. *For any integers $n, N > 1$ there exists a constant $c(n, N) > 0$ such that, for any prime $l > c(n, N)$ and any two rings of integers $\mathcal{O}$ and $\mathcal{O}'$ of finite extensions of $\mathbf{Q}_l$,*

(a) *every abstract homomorphism $\varphi : SL_n(\mathcal{O}') \rightarrow GL_N(\mathcal{O})$ is continuous and $\ker \varphi$ is contained in the subgroup of scalar matrices; and*

(b) *if $N < n$, then there is no non-trivial abstract homomorphism $\varphi : SL_n(\mathcal{O}') \rightarrow GL_N(\mathcal{O})$. Part (a) remains true if we replace SL_n by $GL_n, Sp_{2n}, PSL_n, PGL_n$, or PSp_{2n} . The same goes for Part (b), if for GL_n and PGL_n we require that φ be non-Abelian; and for Sp_{2n} and PSp_{2n} , that $N < 2n$.*

2. MODULI OF ABELIAN VARIETIES

In this section we recall results of Shimura on Abelian varieties with prescribed endomorphism structures. Let A be an Abelian variety over $\mathbf{C}$. Let E be a semisimple algebra over a field k . Let $\theta : E \rightarrow \text{End}_{\mathbf{C}}(A) \otimes_{\mathbf{Z}} \mathbf{Q}$ be an isomorphism, and let $\mathcal{C}$ be a polarization of A . Two such triples $(A, \mathcal{C}, \theta)$ and $(A', \mathcal{C}', \theta')$ are said to be isomorphic to each other if there exists an isomorphism $A \xrightarrow{\lambda} A'$ such that $\lambda(X') \in \mathcal{C}$ for every divisor $X' \in \mathcal{C}'$, and that $\lambda\theta(\alpha) = \theta'(\alpha)\lambda$ for every $\alpha \in E$.

Proposition 1 (Shimura [21, Prop. 5]). *With $\mathcal{P} = (A, \mathcal{C}, \theta)$ as before, suppose further that A is defined over a number field. Then there exists a unique subfield $k/\mathbf{Q}$, called the field of moduli of $(A, \mathcal{C}, \theta)$, which is characterized by the following property: For every automorphism σ of $\mathbf{C}$, the triple $\mathcal{P}^\sigma$ is isomorphic to $\mathcal{P}$ if and only if $\sigma|_k = \text{id}$.*

□

Let F be a totally real number field contained in the center of E . Let ρ be an involution of E/k such that $a^\rho = a$ for every $a \in F$, and that $\text{tr}_{E/\mathbf{Q}}(aa^\rho) > 0$ for every non-zero $a \in E$. Let $n = \dim A$, and let Φ be a representation of L by the $n \times n$ complex matrices containing no zero representations. Following Shimura [23, §1.4], the triple $(A, \mathcal{C}, \theta)$ is called a *polarized Abelian variety of type $\{E, \Phi, \rho\}$* if

- (i) Φ takes the identity element of E to the identity matrix.
- (ii) the representation of $\theta(x)$ for $x \in E$ with respect to any analytic coordinates given by $A \simeq \mathbf{C}^n/(\text{lattice})$ is equivalent to Φ .
- (iii) the involution of $\text{End}_{\mathbf{C}}(A) \otimes \mathbf{Q}$ determined by $\mathcal{C}$ coincides on $\theta(E)$ with the involution $\theta(a) \rightarrow \theta(a^\rho)$.

Proposition 2 (Shimura [23, Prop. 1.6]). *Let $(A, \mathcal{C}, \theta)$ be a polarized Abelian variety of type $\{E, \Phi, \rho\}$. Then the field of moduli of the triple $(A, \mathcal{C}, \theta)$ contains the number field generated over $\mathbf{Q}$ by the trace of the elements $\Phi(\alpha)$ for all α in the center of E .*

□

Next, we recall Albert’s work on the classification of endomorphism algebras of simple complex Abelian varieties. Let A be a simple, polarized Abelian variety over the complex numbers. Then $E := \text{End}(A) \otimes \mathbf{Q}$ is a finite dimensional division algebra over $\mathbf{Q}$ equipped

with the Rosati involution with respect to the polarization. Denote by K the center of E , and by K_0 the fixed field of the Rosati involution restricted to K . Set

$$[E : K] = \delta^2, [K : \mathbf{Q}] = e, [K_0 = \mathbf{Q}] = e_0.$$

It is known that $\delta^2 e$ divides $2 \dim A$, and if $e = 2 \dim A$, then K is a CM field [24, §2.2].

E is said to be a division algebra of the first kind (resp. second kind) if the Rosati involution acts trivially on K (resp. non-trivially). By the work of Albert, E falls into one of four types, exactly three of which correspond to division algebras of the first kind. Moreover, we have the following restrictions on e, e_0 and δ ([14, Prop. 5.5.7]):

$\text{End}(A) \otimes \mathbf{Q}$	δ	e_0	restriction
totally real number field	1	e	$e \mid \dim A$
totally indefinite quaternion algebra	2	e	$2e \mid \dim A$
totally definite quaternion algebra	2	e	$2e \mid \dim A$
division algebra of the second kind	δ	$e/2$	$e_0 \delta^2 \mid \dim A$

TABLE 1.

3. ENDOMORPHISM ALGEBRAS OVER TOTALLY REAL FIELDS

Let $(A, \mathcal{C}, \theta)$ be an absolutely simple, polarized Abelian variety of type $\{E, \Phi, \rho\}$ over k and of *odd* dimension d . Cf. Table 1 and we see that $E = \text{End}_{\bar{k}}(A) \otimes \mathbf{Q}$ is either a totally real number field of degree dividing d or a division algebra of the second kind. To prove Theorem 1 we need to eliminate this second possibility when k is totally real.

Suppose E is a division algebra of the second kind. Following the notation in Table I, we have [22, §2]

$$(3) \quad E \otimes_{\mathbf{Q}} \mathbf{R} \simeq \underbrace{M_{\delta}(\mathbf{C}) \oplus \cdots \oplus M_{\delta}(\mathbf{C})}_{e_0},$$

where $M_{\delta}(\mathbf{C})$ denotes the algebra of $\delta \times \delta$ complex matrices. For $i = 1, \dots, e_0$, denote by $\chi_i : E \otimes_{\mathbf{Q}} \mathbf{R} \rightarrow M_{\delta}(\mathbf{C})$ the projection onto the i -th factor. According to [22, §2], the inequivalent absolutely irreducible representations of E are given by

$$\chi_1, \bar{\chi}_1, \dots, \chi_g, \bar{\chi}_g,$$

where $\bar{\chi}_i$ denotes the complex conjugate of χ_i . Moreover, for each i there exist integers $r_i, s_i \geq 0$ satisfying

$$(4) \quad \begin{aligned} r_i + s_i &= 2\delta(\dim A)/[E : \mathbf{Q}] \\ &= (\dim A)/(\delta e_0) \quad \text{since } [E : \mathbf{Q}] = [E : K][K : K_0][K_0 : \mathbf{Q}] \\ &= \delta^2 \cdot 2 \cdot e_0, \end{aligned}$$

such that with

$$(5) \quad \Phi_i := (\chi_i \otimes I_{r_i}) \oplus (\bar{\chi}_i \otimes I_{s_i})$$

(I_j denotes the $j \times j$ identity matrix), the type $\{E, \Phi, \rho\}$ of A/k satisfies [22, §2],

$$(6) \quad \Phi \simeq \Phi_1 \oplus \cdots \oplus \Phi_g.$$

Apply this to the case where $\dim A$ is odd, it then follows from (4) that $r_i + s_i$ is odd. In particular, $r_i \neq s_i$ for every i .

In light of the isomorphism (3), we can find $\epsilon \in E$ such that

- (i) $\text{trace}(\chi_1(\epsilon))$ is complex, and
- (ii) the imaginary part of $\chi_1(\epsilon)$ is large, but $|\text{trace}(\chi_i(\epsilon))|$ is small for every $i > 1$.

Then

$$\begin{aligned} \text{trace}(\Phi(\epsilon)) &= \sum_i \text{trace}(\Phi_i(\epsilon)) && \text{by (6)} \\ (7) \quad &= \sum_i [r_i \cdot \text{trace}(\chi_i(\epsilon)) + s_i \cdot \overline{\text{trace}(\chi_i(\epsilon))}] && \text{by (5)}. \end{aligned}$$

Since $r_1 \neq s_1$, condition (i) implies that the term $i = 1$ in (7) is not real. Condition (ii) then implies that $\text{trace}(\Phi(\epsilon))$ is not real. By Proposition 2, the field of moduli of $(A, \mathcal{C}, \theta)$ cannot be totally real. On the other hand, by Proposition 1 this field of moduli is contained in k , which is totally real by hypothesis. This is a contradiction, and Theorem 1 follows.

4. ‘ABSTRACT’ HOMOMORPHISMS OF l -ADIC GROUPS

Let G be a linear algebraic group (i.e. a closed subgroup of GL_n) defined over a field K . Fix an algebraic closure $\overline{K}$ of K . The group G is said to be almost simple (resp. absolutely almost simple) if it has no proper, normal subgroup over K (resp. over $\overline{K}$) of dimension > 0 . It is said to be isotropic over K if it contains a non-zero, split K -torus. Both $SL_n(K)$ and $Sp_{2n}(K)$ are isotropic and are absolutely almost simple over $\overline{K}$.

Let $\varphi : K \rightarrow K'$ be a non-trivial field homomorphism. Denote by G_φ the base change of G to K' via φ , i.e. the pullback of $G \rightarrow \text{Spec } K$ via φ :

$$\begin{array}{ccc} G_\varphi & \longrightarrow & G \\ \downarrow & & \downarrow \\ \text{Spec } K' & \longrightarrow & \text{Spec } K. \end{array}$$

If $\rho : G \rightarrow PGL_n(K')$ is a rational representation, i.e. a morphism of algebraic groups, then we get a natural map $\rho_\varphi : G_\varphi \rightarrow PGL_n(K')$.

The following special case of [6, Thm. 10.3] contains all the ingredients from the work of Borel-Tits for our subsequent applications.

Theorem 5 (Borel-Tits). *Let K be a field of characteristics zero, and let K' be an algebraically closed field. Let G be a connected linear algebraic group over K which is absolutely almost simple and is isotropic over K . Let $\rho : G(K) \rightarrow PGL_n(K')$ be an irreducible projective representation ($n \geq 2$). Then there exists finitely many distinct, non-trivial field homomorphisms $\varphi_i : K \rightarrow K'$, and for each i , a non-trivial, irreducible rational projective representation $\pi(i)$ of G , such that*

$$\rho \simeq \otimes_i (\pi(i)_{\varphi_i} \circ \varphi_i).$$

In particular, $\text{char}(K') = 0$. The pairs consisting of φ_i and the equivalence class of π_i are unique up to reordering.

□

The following fact is elementary.

Lemma 1. *Let $1 \rightarrow C \rightarrow G \rightarrow \tilde{G} \rightarrow 1$ be a central extension of groups. Given any group H , any two lifts of a homomorphism $H \rightarrow \tilde{G}$ differs by the multiple of a homomorphism $H \rightarrow C$.* $\square$

For the rest of this section, K denotes a field of characteristic zero. Let $i \neq j$ be integer between 1 and n . For any elements $\alpha \in K^\times$ and $\beta \in K$, define two matrices

$$\begin{aligned} A_i(\alpha) &= \text{the matrix obtained from the } 2 \times 2 \text{ identity matrix by replacing the} \\ &\quad ii\text{-th entry by } \alpha, \\ B_{ij}(\beta) &= \text{the matrix obtained from the } 2 \times 2 \text{ identity matrix by adding } \beta \text{ to the} \\ &\quad ij\text{-th entry.} \end{aligned}$$

For $i \neq j$ we have the identity

$$(8) \quad A_i(\alpha) B_{ij}(\beta) A_i(\alpha)^{-1} = B_{ij}(\alpha^{\text{sgn}(j-i)} \beta).$$

Proposition 3. *Every abstract, absolutely irreducible representation $GL_2(K) \rightarrow GL_d(\overline{K})$ is equivalent to*

$$[\otimes_{s=1}^r (\sigma_{\varphi_s}^{(n_s)} \circ \varphi_s)] \otimes (f \circ \det),$$

where V is the underlying representation space associated to natural representation of $GL_2(K)$; $\det : GL_2(K) \rightarrow K^\times$ is the determinant map; and $f : K^\times \rightarrow \overline{K}^\times$ is a homomorphism of multiplicative groups.

Proof. Since ρ is absolutely irreducible, so are $\rho|_{SL_2(K)}$ and

$$\tilde{\rho}_S := \text{the projective representation } SL_2(K) \rightarrow PGL_d(\overline{K}) \text{ associated to } \rho|_{SL_2(K)}.$$

Define

$$\widetilde{\sigma_{\varphi_s}^{(n_s)}} := \text{the projective representation associated to } \sigma_{\varphi_s}^{(n_s)}.$$

The absolutely irreducible rational projective representations of $SL_2(K)$ are precisely the restrictions to $SL_2(K)$ of the $\widetilde{\sigma_{\varphi_s}^{(n_s)}}$, so by Theorem 5,

$$\tilde{\rho}_S \simeq \otimes_{s=1}^r [(\widetilde{\sigma_{\varphi_s}^{(n_s)}} \circ \varphi_s)|_{SL_2(K)}],$$

where $n_s \geq 1$ and the $\varphi_s : K \rightarrow \overline{K}$ are field homomorphisms. Since $SL_2(K)$ has no non-trivial Abelian quotients, by Lemma 1 there is a unique lift of $\tilde{\rho}_S$ to $\rho|_{SL_2(K)}$, whence

$$(9) \quad \rho|_{SL_2(K)} \simeq \otimes_{s=1}^r [(\sigma_{\varphi_s}^{(n_s)} \circ \varphi_s)|_{SL_2(K)}].$$

To determine ρ it then remains to study its action on the matrices $A_i(\alpha)$ with $\alpha \in K^\times$.

The identity (8) gives

$$\sigma_{\varphi}^{(n)} B_{ij}(\varphi(\alpha^{\text{sgn}(j-i)} \beta)) = [\sigma_{\varphi}^{(n)} A_i(\varphi(\alpha))] [\sigma_{\varphi}^{(n)} B_{ij}(\varphi(\beta))] [\sigma_{\varphi}^{(n)} A_i(\varphi(\alpha))]^{-1},$$

whence

$$\begin{aligned} & [\otimes_{s=1}^r \sigma_{\varphi_s}^{(n_s)} B_{ij}(\varphi_s(\alpha^{\text{sgn}(j-i)} \beta))] \\ &= [\otimes_{s=1}^r \sigma_{\varphi_s}^{(n_s)} A_i(\varphi_s(\alpha))] [\otimes_{s=1}^r \sigma_{\varphi_s}^{(n_s)} B_{ij}(\varphi_s(\beta))] [\otimes_{s=1}^r \sigma_{\varphi_s}^{(n_s)} A_i(\varphi_s(\alpha))]^{-1}. \end{aligned}$$

In light of (9), this relation holds if we replace each $\otimes_{s=1}^r \sigma_{\varphi_s}^{(n_s)} B_{ij}(\varphi_s(-))$ by $\rho(B_{ij}(-))$. It follows that

$$\left[\otimes_{s=1}^r \sigma_{\varphi_s}^{(n_s)} A_i(\varphi_s(\alpha)) \right] \left[\rho(A_i(\alpha)) \right]^{-1}$$

commutes with every $B_{ij}(\varphi_s(\gamma))$. As γ runs through all elements of K , the matrices $B_{ij}(\varphi_s(\gamma))$ generate $SL_2(\varphi_s(K))$ [4, Thm. 4.6]. Since $\rho|_{SL_2(K)}$ is absolutely irreducible, Schur’s Lemma implies that

$$\rho(A_i(\alpha)) = \left[\otimes_{s=1}^r \sigma_{\varphi_s}^{(n_s)} A_i(\varphi_s(\alpha)) \right] f(\alpha)$$

for some element $\alpha \in \overline{K}^\times$. Since ρ is a homomorphism, $f : K^\times \rightarrow \overline{K}^\times$ is in fact a homomorphism of multiplicative groups. Thus

$$\rho = \left[\otimes_{s=1}^r (\sigma_{\varphi_s}^{(n_s)} \circ \varphi_s) \right] \otimes (f \circ \det)$$

when restricted to $SL_2(K)$ and to the subgroup of $GL_2(K)$ generated by $A_i(\alpha)$ for all the $\alpha \in K^\times$. These two subgroups together generate $GL_2(K)$, so we are done. $\square$

For every $s \in SL_2(K)$, the trace of $\sigma^{(n)}(s)$ is a $\mathbf{Z}$ -polynomial of degree n in $\text{trace}(s)$. Write

$$T_n(\text{trace}(s))$$

for this polynomial.

Lemma 2. *Suppose there exists an infinite subset $\Sigma \subset SL_2(K)$ of elements with pairwise distinct traces, such that*

$$(10) \quad \prod_{i=1}^r T_{n_i}(\text{trace}(s)) = \prod_{j=1}^t T_{m_j}(\text{trace}(s))$$

for every $s \in \Sigma$. Then $r = t$ and, up to reordering, $n_i = m_i$ for all i .

Proof. Denote by α and $1/\alpha$ the eigenvalues in $\overline{K}$ of $s \in \Sigma$. Write ζ_m for a fixed primitive m -th root of unity. Then

$$T_n(\text{trace}(s)) = \frac{\alpha^{n+1} - (1/\alpha)^{n+1}}{\alpha - (1/\alpha)} = \prod_{u=1}^n (\alpha - \zeta_{n+1}^u (1/\alpha)).$$

The equality (10) then becomes

$$\prod_{i=1}^r \prod_{u=1}^{n_i} (\alpha - \zeta_{1+n_i}^u (1/\alpha)) = \prod_{j=1}^t \prod_{v=1}^{m_j} (\alpha - \zeta_{1+m_j}^v (1/\alpha)).$$

Since this holds for infinitely many distinct $\alpha \in \overline{K}$, this becomes an equality of monic Laurent polynomials in the variable α over the field $L = K(\{\zeta_{1+n_i}, \zeta_{1+m_j}\}_{i,j})$. Since the ring of (one variable) Laurent polynomials over L has unique factorization, we are done. $\square$

5. ODD-DIMENSIONAL l -ADIC REPRESENTATIONS

Proof of Theorem 2. Suppose E is a totally real field. We consider two cases.

Case: $[E : \mathbf{Q}] = 1$, so $\text{End}_{\bar{k}}(A) = \mathbf{Z}$.

Since d is odd, Serre [20, 2.2.7 – 2.2.8] shows that the image of $\rho_{A,l}$ contains $Sp_{2d}(\mathbf{Z}_l)$ for all but finitely many l . Following the notation in (1), the restriction of the abstract homomorphism ι_λ to $Sp_{2d}(\mathbf{Q}_l)$ then gives rise to a non-trivial homomorphism $\iota'_\lambda : Sp_{2d}(\mathbf{Z}_l) \rightarrow GL_d(F_\lambda)$ for some finite extension $F_\lambda/\mathbf{Q}_l$. Since $Sp_{2d}(\mathbf{Z}_l)$ is compact, the argument in [18, Remark 1 on p. 1] shows that ι'_λ is equivalent to a homomorphism with image in $GL_d(\mathcal{O}_{F_\lambda})$. By Theorem 4(b), this is impossible for l sufficiently large.

Case: $[E : \mathbf{Q}] = d > 1$.

Suppose $\rho_\lambda : G_k \rightarrow GL_d(F_\lambda)$ is an abstract, absolutely irreducible representation coming from A/k . Following the notation in (1) – (2), that means ρ_λ is the composition of some $\tau_{l,m} : G_k \rightarrow GL_2(E_{l,m})$ with an abstract homomorphism $\psi : GL_2(E_{l,m}) \rightarrow GL_d(\overline{\mathbf{Q}}_l)$. Since ρ_λ is absolutely irreducible, so does ψ . Apply Proposition 3 and we are done. $\square$

Proof of Theorem 3. Our goal is to understand the dependency of the parameters $r(\lambda)$, $n_i(\lambda)$, f_λ and $\varphi_i(\lambda)$ on λ . Faltings shows that $\tau_{l,m}$ is semisimple, and that $E_{l,m}$ is the commutant of $\tau_{l,m}(G_k)$ in the $\text{End}(T_l(A))$. The argument in ([16, Thm. 4.5.4]) then shows that the Lie algebra of $\tau_{l,m}(G_k)$ is equal to $\{u \in \mathfrak{gl}_2(E_{l,m}) : \text{trace}(u) \in \mathbf{Q}_l\}$. Since $\tau_{l,m}$ is equivalent to a representation whose image lies in $GL_2(\mathcal{O}_{E_{l,m}})$ (cf. [18, Remark 1 on p. 1]), it follows that

$$(11) \quad \begin{aligned} & \text{for every } l, \tau_{l,m}(G_k) \text{ is (conjugate to) an open} \\ & \text{subgroup in } \{g \in GL_2(\mathcal{O}_{E_{l,m}}) : \det g \in \mathbf{Z}_l^\times\}. \end{aligned}$$

In particular, there exists an infinite subset $\{g_i\}_i \subset G_k$ such that, for a fixed l , $\tau_{l,m}(g_i) \in SL_2(\mathbf{Z}_l) \subset SL_2(\mathcal{O}_{E_{l,m}})$ and with pairwise distinct traces. Since $\{\tau_{l,m}\}_l$ is a strictly compatible family, that means $\tau_i := \text{trace}(\tau_{l,m}(g_i))$ lies³ in $\mathbf{Q}$ and is independent of l .

Let $\lambda, \mathfrak{q}$ be two distinct finite places of F of residual characteristic l and q ($l = q$ is allowed). Following the notation in Lemma 2, we have

$$\begin{aligned} \Phi_\lambda^{-1} \left(\prod_{i=1}^{r(\lambda)} T_{n_i(\lambda)}(\tau_i) \right) &= \Phi_\lambda^{-1} \left(\text{trace}(\rho_\lambda(g_i)) \right) \\ &= \Phi_{\mathfrak{q}}^{-1} \left(\text{trace}(\rho_{\mathfrak{q}}(g_i)) \right) \quad \begin{array}{l} \{\rho_\lambda\}_\lambda \text{ is a strictly compatible family} \\ \text{and the Frobenius are dense in } G_k \end{array} \\ &= \Phi_{\mathfrak{q}}^{-1} \left(\prod_{j=1}^{r(\mathfrak{q})} T_{n_j(\mathfrak{q})}(\tau_i) \right) \end{aligned}$$

Since this holds for all τ_i , by Lemma 2 that means $r(\lambda) = r(\mathfrak{q})$ and, up to reordering, $n_i(\lambda) = n_i(\mathfrak{q})$ for all i .

It remains to study the multiplicative homomorphism $f_\lambda : E_l^\times \rightarrow \mathbf{Q}_l^\times$. Since $\det \tau_{l,m}$ is the l -adic cyclotomic character [16], it is surjective onto $\mathbf{Z}_l^\times$ for all but finitely many l . Since $\mathbf{Z}$ is dense in $\mathbf{Z}_l$, we can find an infinite subset $\{h_i\}_i \subset G_k$ and a prime l such

³in fact these traces lie in $\mathbf{Z}$, although we do not need it.

that $\{\det \pi_{l,m}(h_i)\}_i = \mathbf{Z} - \{0\}$. Moreover, $d_i := \det \pi_{l,m}(h_i)$ is independent of l . From $\text{trace}(\rho_\lambda(h_i)) = \text{trace}(\rho_q(h_i))$ for all i we then see that $f_\lambda(d_i) = f_q(d_i)$ for all i , whence $f_\lambda|_{\mathbf{Z}} = f_q|_{\mathbf{Z}}$. $\square$

6. CONTINUITY OF l -ADIC MAPS

Lemma 3. *For any $n > 1$ there exists a constant $c_1(n) > 0$, such that the degree of any non-Abelian, complex irreducible representation of $SL_n(\mathbf{Z}/l)$, $GL_n(\mathbf{Z}/l)$, $Sp_{2n}(\mathbf{Z}/l)$, $PSL_n(\mathbf{Z}/l)$, $PGL_n(\mathbf{Z}/l)$, and $PSp_{2n}(\mathbf{Z}/l)$ is $> c_1(n)l$.*

Proof. Every irreducible representation of $PSL_n(\mathbf{Z}/l)$, $PGL_n(\mathbf{Z}/l)$ and $PSp_{2n}(\mathbf{Z}/l)$ gives rise to an irreducible representation of $SL_n(\mathbf{Z}/l)$, $GL_n(\mathbf{Z}/l)$ and $Sp_{2n}(\mathbf{Z}/l)$, respectively, so it suffices to examine the three latter class of groups.

By Schur’s Lemma, any irreducible complex representation $\rho : SL_n(\mathbf{Z}/l) \rightarrow GL(V)$ takes the center of $SL_n(\mathbf{Z}/l)$ to the center of $GL(V)$. Compose ρ with the canonical map $SL(V) \rightarrow PGL(V)$ and we get an irreducible *projective* representation $PSL_n(\mathbf{Z}/l) \rightarrow PGL(V)$. To bound the minimal degree of non-Abelian irreducible representations of $SL_n(\mathbf{Z}/l)$, it then suffices to bound the minimal degree of non-Abelian irreducible *projective* representations of $PSL_n(\mathbf{Z}/l)$. This is done in [13], and the lower bound we desire follows from the main theorem there. The same discussion applies to Sp_{2n} as well.

Finally, let χ be an irreducible representation of GL_n . Its restriction to SL_n remains irreducible [25, Lem. 4.5], so $\deg \chi > 1$ if χ is non-Abelian since SL_n is its own commutator subgroup. The Lemma for GL_n is then reduced to the case of SL_n . This completes the proof of the Lemma. $\square$

Lemma 4. *Fix integers $m, N \geq 1$ and $n \geq 2$. Then there exists a constant $c_2(m, n, N) > 0$ such that, for any prime $l > c_2(m, n, N)$, no subgroup of $GL_N(\mathbf{C})$ is isomorphic to $SL_n(\mathbf{Z}/l^m)$, $GL_n(\mathbf{Z}/l^m)$, $Sp_{2n}(\mathbf{Z}/l^m)$, $PSL_n(\mathbf{Z}/l^m)$, $PGL_n(\mathbf{Z}/l^m)$, or $PSp_{2n}(\mathbf{Z}/l^m)$.*

Proof. Suppose there exists an injective map $\phi : SL_n(\mathbf{Z}/l^m) \rightarrow GL_N(\mathbf{C})$. We can assume that ϕ is irreducible. Denote by S the kernel of the reduction map $SL_n(\mathbf{Z}/l^m) \rightarrow SL_n(\mathbf{Z}/l)$. Every irreducible representation of a nilpotent group P is induced from a degree one character of some subgroup of P [9, Thm. 52.1], so the degree of every irreducible representation of the l -group S is either 1 or is divisible by l . But $\deg(\phi|_S) = N$, so for $l > N$ every irreducible representation of S appearing in $\phi|_S$ has degree 1. Thus $\phi(S)$ is Abelian. But S is non-Abelian if $m > 2$, so $m \leq 2$. Lemma 3 covers the case $m = 1$, so it remains to consider the case $m = 2$. We first introduce some notation.

Let H be a normal subgroup of a finite group G . For any character ψ and any element $g \in G$, denote by ψ^g the *conjugate* of ψ by g :

$$(12) \quad \psi^g(h) := \psi(g^{-1}hg) \quad \text{for all } h \in H.$$

We now resume the proof of the Lemma. Take $m = 2$, so S is Abelian. Clifford’s theory [9, §49] gives a decomposition

$$(13) \quad \phi|_S \simeq (\psi^{s_1} \oplus \cdots \oplus \psi^{s_r})^e,$$

where $e, r \geq 1$ are integers, $s_i \in SL_n(\mathbf{Z}/l^2)$, and ψ is an irreducible representation of S . Moreover, $SL_n(\mathbf{Z}/l^2)$ permutes the ψ^{g_i} transitively via (12). We claim that this action on the ψ^{g_i} , when restricted to S , is trivial. To see this, note that for any $h, s \in S$,

$$\begin{aligned} (\psi^{g_i})^s(h) &= \psi(s^{-1}g_i^{-1}hg_is) \\ &= \psi(g_i^{-1}\sigma^{-1}h\sigma g_i) \quad \text{for some } \sigma \in S, \text{ since } S \text{ is normal} \\ &= \psi(g_i^{-1}hg_i) \quad \text{since } S \text{ is Abelian} \\ &= \psi^{g_i}(h). \end{aligned}$$

This verifies the claim. As a result, $SL_n(\mathbf{Z}/l^2)/S \simeq SL_n(\mathbf{Z}/l)$ permutes the ψ^{g_i} transitively. The left side of (13) has degree N , so $r \leq N$. Thus $SL_n(\mathbf{Z}/l)$ permutes transitively a set of size $r \leq N$. We have two cases:

- $r = 1$:
Then $\phi|_S$ is the direct sum of copies of a fixed irreducible representation ψ of S . Since S is Abelian, that means $\deg \psi = 1$, whence $\phi|_S$ has cyclic image. This is impossible since S is non-cyclic and ϕ is injective.
- $1 < r \leq N$:
Then $SL_n(\mathbf{Z}/l)$ contains a proper subgroup of index $\leq N$. That means $SL_n(\mathbf{Z}/l)$ has a non-trivial permutation representation, and hence a non-trivial irreducible representation, of degree $\leq N$. This is impossible if l is large enough, by Lemma 3 (note that $SL_n(\mathbf{Z}/l)$ has no non-trivial degree one character if $(n, l) \neq (2, 2)$).

This completes the proof of the Lemma for SL_n . The same argument applies to PSL_n, Sp_{2n} and PSp_{2n} . Finally, if $GL_N(\mathbf{C})$ contains a subgroup isomorphic to $GL_n(\mathbf{Z}/l^m)$ or $PGL_n(\mathbf{Z}/l^m)$, then it contains one isomorphic to $SL_n(\mathbf{Z}/l^m)$ or $PSL_n(\mathbf{Z}/l^m)$, so we are done. $\square$

Proposition 4. *Let A be a local ring with maximal ideal λ . Suppose $\text{char}(A/\lambda) > 3$. Then a subgroup $H \subset GL_N(A)$ is normal if and only if there exists an ideal $\mathfrak{a} \subset A$ such that the image of H in $GL_N(A/\mathfrak{a})$ consists of the scalar matrices. The same holds for $SL_N, Sp_{2N}, PGL_N, PSL_N$ and PSp_{2N} .*

Proof. For $GL_N(A)$ and $SL_N(A)$ see [15, p. 84 and p. 245]. For $Sp_{2d}(A)$ see [15, p. 210]. $\square$

Proof of Theorem 4. Again we give the argument for SL_n . We can assume that $l > 3$. Denote by λ and λ' the maximal ideal of $\mathcal{O}$ and $\mathcal{O}'$, respectively.

(a) We can assume that φ is not trivial. For $i > 0$, denote by G_i the kernel of the reduction map $GL_N(\mathcal{O}) \rightarrow GL_N(\mathcal{O}/\lambda^i)$. Every quotient G_i/G_{i+1} is a finite Abelian l -group, whence the same holds for $\varphi^{-1}(G_i)/\varphi^{-1}(G_{i+1})$.

Lemma 5. *Every $\varphi^{-1}(G_i)$ has finite index in $SL_n(\mathcal{O}')$.*

Proof. Every G_i is normal in $GL_N(\mathcal{O})$, so every $\varphi^{-1}(G_i)$ is normal in $SL_n(\mathcal{O}')$. In light of Proposition 4, it suffices to show that none of the $\varphi^{-1}(G_i)$ is contained in the center of $SL_n(\mathcal{O}')$. Suppose otherwise. Then $SL_n(\mathcal{O}')/\varphi^{-1}(G_i)$ is infinite and injects into $GL_N(\mathcal{O})/G_i \simeq GL_N(\mathcal{O}/\lambda^i)$, which is finite. This is a contradiction. $\square$

The G_i form a basis of open neighborhoods of the identity element in $GL_N(\mathcal{O})$. Lemma 5 then implies that $\varphi^{-1}(U)$ has finite index in $SL_n(\mathcal{O}')$ for every sufficiently small open set $U \subset GL_N(\mathcal{O})$ containing the identity; and hence for all open sets $U \subset GL_N(\mathcal{O})$. That means φ is continuous.

Finally, suppose $\ker \varphi$ is not contained in the subgroup of scalar matrices. In light of Proposition 4, for some $m \geq 1$ we then get an injective map of $SL_n(\mathcal{O}'/\lambda'^m)$ or $PSL_n(\mathcal{O}'/\lambda'^m)$ into $GL_N(\mathcal{O})$. Restrict this map to the subgroup $SL_n(\mathbf{Z}/l^m)$ or $PSL_n(\mathbf{Z}/l^m)$ and note that we have a (discontinuous) inclusion $GL_N(\mathcal{O}) \subset GL_N(\mathbf{C})$, we get a contradiction, by Lemma 4. This completes the proof of Part (a).

(b) We claim that if such φ exists, then we can find an injective map of abstract groups

$$SL_n(\mathbf{F}_l) \hookrightarrow GL_N(\overline{\mathbf{F}}_l) \quad \text{or} \quad PSL_n(\mathbf{F}_l) \hookrightarrow GL_N(\overline{\mathbf{F}}_l).$$

By the work of Steinberg [26] it then follows that $N \geq n$.

For every integer $i \geq 1$, denote by G_i the kernel of the reduction map $GL_N(\mathcal{O}) \rightarrow GL_N(\mathcal{O}/\lambda^i)$. Every quotient G_i/G_{i+1} is a finite Abelian l -group, whence the same holds for the $\varphi^{-1}(G_i)/\varphi^{-1}(G_{i+1})$. We claim that

$$(14) \quad \varphi^{-1}(G_1) \neq SL_n(\mathcal{O}').$$

Suppose otherwise. Since $\ker(\varphi)$ is a proper subgroup of $SL_n(\mathcal{O}')$ and since $\cap_i G_i = \{I\}$, if (14) is false then there exists a smallest integer $j \geq 1$ such that $\varphi^{-1}(G_{j+1}) \subsetneq \varphi^{-1}(G_j)$; necessarily $\varphi^{-1}(G_j) = SL_n(\mathcal{O}')$. On the other hand, every $\varphi^{-1}(G_i)$ is normal in $SL_n(\mathcal{O}')$. Since $l > 3$, Proposition 4 implies that

$$\varphi^{-1}(G_{j+1}) = \ker(SL_n(\mathcal{O}') \xrightarrow{\text{mod } \lambda'^m} SL_n(\mathcal{O}'/\lambda'^m))$$

or

$$\varphi^{-1}(G_{j+1}) = \ker(SL_n(\mathcal{O}') \xrightarrow{\text{mod } \lambda'^m} PSL_n(\mathcal{O}'/\lambda'^m))$$

for some $m \geq 1$. Then $\varphi^{-1}(G_j)/\varphi^{-1}(G_{j+1}) = SL_n(\mathcal{O}')/\varphi^{-1}(G_{j+1}) \simeq SL_n(\mathcal{O}'/\lambda'^m)$ or $PSL_n(\mathcal{O}'/\lambda'^m)$, which is not a finite Abelian l -group. This is a contradiction, so (14) must hold, and we get an injective map

$$SL_n(\mathcal{O}')/\varphi^{-1}(G_1) \hookrightarrow GL_N(\mathcal{O}_L)/G_1 \simeq GL_N(\mathcal{O}_L/\lambda).$$

By Proposition 4, we get an injective map

$$(15) \quad SL_n(\mathcal{O}'/\lambda'^m) \hookrightarrow GL_N(\mathcal{O}_L/\lambda) \quad \text{or} \quad PSL_n(\mathcal{O}'/\lambda'^m) \hookrightarrow GL_N(\mathcal{O}_L/\lambda).$$

for some $m \geq 1$. Every element of $GL_N(\mathcal{O}_L/\lambda)$ either is semisimple or contains a non-trivial Jordan block (over $\overline{\mathbf{F}}_l$). In the first case the element has order prime to l . In the second case, if $l > N$ then every non-semisimple element has order divisible by l but not by l^2 . On the other hand, since $l > 2$, both $SL_n(\mathcal{O}'/\lambda'^m)$ and $PSL_n(\mathcal{O}'/\lambda'^m)$ contain elements of order l^m (coming from transvections). Since the maps in (15) are injective, that means $m = 1$. This completes the proof of Part (b). □

ACKNOWLEDGEMENT

My interest in odd-dimensional Galois representations was inspired by a lecture of Ash on the analog of Serre's conjecture for $GL(3)$ modular representations. I am indebted to Humphreys for his help with the work of Borel-Tits, and to Ribet for his help concerning the image of λ -adic representations. I would like to thank Dettweiler for pointing out a mistake in an earlier draft. I also benefited from a useful conversation with Rosen.

REFERENCES

- [1] G. Allison, A. Ash and E. Avner, Galois representations, Hecke operators, and the mod- p cohomology of $GL(3, \mathbf{Z})$ with twisted coefficients. *Experiment. Math.* **7** (1998) 361-390.
- [2] A. Ash, D. Grayson and P. Green, Computations of cuspidal cohomology of congruence subgroups of $SL(3, \mathbf{Z})$. *J. Number Theory* **19** (1984) 412-436.
- [3] A. Ash, R. Pinch and R. Taylor, An $\widehat{A_4}$ extension of $\mathbf{Q}$ attached to a nonselfdual automorphic form on $GL(3)$. *Math. Ann.* **291** (1991) 753-766.
- [4] E. Artin, *Geometric algebra*. Interscience Publ., 1957.
- [5] D. Blasius and J. Rogawski, Galois representations for Hilbert modular forms. *Bull. Amer. Math. Soc. (N.S.)* **21** (1989) 65-69.
- [6] A. Borel and J. Tits, Homomorphismes 'abstraits' de groupes algébriques simples. *Ann. Math. (2)* **97** (1973) 499-571.
- [7] W. C. Chi, l -adic and λ -adic representations associated to abelian varieties defined over number fields. *Amer. J. Math.* **114** (1992) 315-353.
- [8] L. Clozel, Motifs et formes automorphes: applications du principe de fonctorialité, in: *Automorphic forms, Shimura varieties, and L-functions*, Vol. I (Ann Arbor, MI, 1988) 77-159. Academic Press, 1990.
- [9] C. W. Curtis and I. Reiner, *Representation theory of finite groups and associative algebras*. Interscience Publ., 1962.
- [10] W. Fulton and J. Harris, *Representation theory*. Springer-Verlag, 1991.
- [11] B. van Geemen and J. Top, A non-selfdual automorphic representation of GL_3 and a Galois representation. *Invent. Math.* **117** (1994) 391-401.
- [12] B. van Geemen and J. Top, Selfdual and non-selfdual 3-dimensional Galois representations. *Compositio Math.* **97** (1995) 51-70.
- [13] V. Landazuri and G. M. Seitz, On the minimal degrees of projective representations of the finite Chevalley groups. *J. Algebra* **32** (1974) 418-443.
- [14] H. Lange and C. Birkenhake, *Complex Abelian varieties*. Springer-Verlag, 1992.
- [15] B. R. McDonald, *Geometric algebra over local rings*. M. Dekker, 1976.
- [16] K. Ribet, Galois action on division points of Abelian varieties with real multiplications. *Amer. J. Math.* **98** (1976) 751-804.
- [17] J. Schwermer, Cohomology of arithmetic groups, automorphic forms and L -functions, in: *Cohomology of arithmetic groups and automorphic forms (Luminy-Marseille, 1989)*, 1-29. LNM 1447, Springer-Verlag, 1990.
- [18] J. P. Serre, *Abelian l -adic representations and elliptic curves*. Benjamin, New York, 1968.
- [19] J. P. Serre, Représentations l -adiques, in: *Algebraic number theory* (Kyoto Internat. Sympos., Res. Inst. Math. Sci., Univ. Kyoto, Kyoto, 1976), 177-193.
- [20] J. P. Serre, Algèbre et géométrie. *Ann. Collège France* **86** (1985/86) 95-100.
- [21] G. Shimura, On the theory of automorphic functions. *Ann. Math. (2)* **70** (1959) 101-144.
- [22] G. Shimura, On analytic families of polarized abelian varieties and automorphic functions. *Ann. Math. (2)* **78** (1963) 149-192.
- [23] G. Shimura, On the field of definition for a field of automorphic functions. *Ann. Math. (2)* **80** (1964) 160-189.
- [24] G. Shimura, *Abelian varieties with complex multiplications and modular functions*. Princeton Univ. Press, 1998.

- [25] W. A. Simpson and J. S. Frame, The character tables for $SL(3, q)$, $SU(3, q^2)$, $PSL(3, q)$, $PSU(3, q^2)$. *Canad. J. Math.* **25** (1973) 486-494.
- [26] R. Steinberg, Representations of algebraic groups. *Nagoya Math. J.* **22** (1963) 33-56.
- [27] R. Taylor, On Galois representations associated to Hilbert modular forms. *Invent. Math.* **98** (1989) 265-280.
- [28] H. Weyl, *Classical groups*, 2nd. ed. Princeton University Press, 1946.

DEPARTMENT OF MATHEMATICS & STATISTICS, UNIVERSITY OF MASSACHUSETTS. AMHERST, MA
01003-4515 USA

E-mail address: `siman@math.umass.edu`